

Tables of prehomogeneous modules and étale modules of reductive algebraic groups

Compiled by Wolfgang Globke,
based on the results cited in the reference section.

Contents

1	Classification of prehomogeneous modules	2
1.1	Irreducible reduced prehomogeneous modules	2
1.2	Non-irreducible simple prehomogeneous modules	4
1.3	Two-simple prehomogeneous modules of type I	7
1.4	Two-simple prehomogeneous modules of type II	12
2	Tables of étale modules	19
2.1	Étale modules with center GL_1	19
2.2	Étale modules for groups with factor Sp_m	19
2.3	One-simple étale modules	19
2.4	Two-simple étale modules	19
	References	24

1 Classification of prehomogeneous modules

1.1 Irreducible reduced prehomogeneous modules

The irreducible and reduced prehomogeneous modules were classified by Sato and Kimura, thus we will label each class by SK n , where n is the number given to the class in §7 of the the original work by Sato and Kimura [8]. Along with each module, we will state the connected component of the generic isotropy group, denoted by G_v° , and in some cases the irreducible relative invariant, denoted by f .

Let G be a reductive group and (G, ϱ, V) an irreducible and reduced prehomogeneous module. Then it is equivalent to one of the following prehomogeneous modules:

SK I Regular irreducible reduced prehomogeneous modules.

1. $(G \times \mathrm{GL}_m, \varrho \otimes \omega_1, V^m \otimes \mathbb{C}^m)$,
where $\varrho : G \rightarrow \mathrm{GL}(V^m)$ is an m -dimensional irreducible representation of a connected semisimple algebraic group G (or $G = \{1\}$ and $m = 1$). We have $G_v^\circ \cong G$ and $f(x) = \det(x)$ for $x \in \mathrm{Mat}_m \cong V^m \otimes \mathbb{C}^m$, $\deg(f) = m$.
2. $(\mathrm{GL}_n, 2\omega_1, \mathrm{Sym}^2 \mathbb{C}^n)$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{SO}_n$ and $f(x) = \det(x)$ for $x \in \{A \in \mathrm{Mat}_n \mid A^\top = A\} \cong \mathrm{Sym}^2 \mathbb{C}^n$, $\deg(f) = n$.
3. $(\mathrm{GL}_{2n}, \omega_2, \bigwedge^2 \mathbb{C}^{2n})$ for $n \geq 3$.
We have $G_v^\circ \cong \mathrm{Sp}_n$ and $f(x) = \mathrm{Pf}(x)$ for $x \in \{A \in \mathrm{Mat}_{2n} \mid A^\top = -A\} \cong \bigwedge^2 \mathbb{C}^{2n}$, $\deg(f) = n$.
4. $(\mathrm{GL}_2, 3\omega_1, \mathrm{Sym}^3 \mathbb{C}^2)$.
We have $G_v^\circ \cong \{1\}$ and $f(a) = a_2^2 a_3^2 + 18a_1 a_2 a_3 a_4 - 4a_1 a_3^3 - 4a_2^3 a_4 - 27a_1^2 a_4^2$ for $a = a_1 x^3 + a_2 x^2 y + a_3 x y^2 + a_4 y^3 \in \mathrm{Sym}^3 \mathbb{C}^2$ (so f is the discriminant of a binary cubic form $a(x, y)$).
5. $(\mathrm{GL}_6, \omega_3, \bigwedge^3 \mathbb{C}^6)$.
We have $G_v^\circ \cong \mathrm{SL}_3 \times \mathrm{SL}_3$ and $f(x) = (x_0 y_0 - \mathrm{trace}(XY))^2 + 4x_0 \det(Y) + 4y_0 \det(X) - 4 \sum_{i,j} \det(X_{ij}) \det(Y_{ji})$ (see §5, p. 83 in [8] for a definition), $\deg(f) = 4$.
6. $(\mathrm{GL}_7, \omega_3, \bigwedge^3 \mathbb{C}^7)$.
We have $G_v^\circ \cong \mathrm{G}_2$ and $\deg(f) = 7$.
7. $(\mathrm{GL}_8, \omega_3, \bigwedge^3 \mathbb{C}^2)$.
We have $G_v^\circ \cong \mathrm{SL}_3$ and $\deg(f) = 16$.
8. $(\mathrm{SL}_3 \times \mathrm{GL}_2, 2\omega_1 \otimes \omega_1, \mathrm{Sym}^2 \mathbb{C}^3 \otimes \mathbb{C}^2)$.
We have $G_v^\circ \cong \{1\}$ and $f(A, B) = \mathrm{dis}(\det(xA + yB))$ for $(A, B) \in \{(X, Y) \mid X, Y \in \mathrm{Mat}_3, X^\top = X, Y^\top = Y\} \cong \mathrm{Sym}^2 \mathbb{C}^3 \otimes \mathbb{C}^2$, $\deg(f) = 12$.
9. $(\mathrm{SL}_6 \times \mathrm{GL}_2, \omega_2 \otimes \omega_1, \bigwedge^2 \mathbb{C}^6 \otimes \mathbb{C}^2)$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SL}_2 \times \mathrm{SL}_2$ and $f(A, B) = \mathrm{dis}(\mathrm{Pf}(xA + yB))$ for $(A, B) \in \{(X, Y) \mid X, Y \in \mathrm{Mat}_6, X^\top = -X, Y^\top = -Y\} \cong \bigwedge^2 \mathbb{C}^6 \otimes \mathbb{C}^2$, $\deg(f) = 12$.
10. $(\mathrm{SL}_5 \times \mathrm{GL}_3, \omega_2 \otimes \omega_1, \bigwedge^2 \mathbb{C}^5 \otimes \mathbb{C}^3)$.
We have $G_v^\circ \cong \mathrm{SL}_2$ and $\deg(f) = 15$.
11. $(\mathrm{SL}_5 \times \mathrm{GL}_4, \omega_2 \otimes \omega_1, \bigwedge^2 \mathbb{C}^5 \otimes \mathbb{C}^4)$.
We have $G_v^\circ \cong \{1\}$ and $\deg(f) = 40$.
12. $(\mathrm{SL}_3 \times \mathrm{SL}_3 \times \mathrm{GL}_2, \omega_1 \otimes \omega_1 \otimes \omega_1, \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^2)$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{GL}_1$ and $f(A, B) = \mathrm{dis}(\det(xA + yB))$ for $(A, B) \in \mathrm{Mat}_3 \oplus \mathrm{Mat}_3 \cong \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^2$, $\deg(f) = 12$.

13. $(\mathrm{Sp}_n \times \mathrm{GL}_{2m}, \omega_1 \otimes \omega_1, \mathbb{C}^{2n} \otimes \mathbb{C}^{2m})$ for $n \geq 2m \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_m \times \mathrm{Sp}_{n-m}$ and $f(X) = \mathrm{Pf}(X^\top JX)$ for $X \in \mathrm{Mat}_{2n, 2m}$, $\deg(f) = 2m$.
14. $(\mathrm{GL}_1 \times \mathrm{Sp}_3, \mu \otimes \omega_3, \mathbb{C} \otimes V^{14})$.
We have $G_v^\circ \cong \mathrm{SL}_3$ and $\deg(f) = 4$, where f is given by the restriction of the relative invariant of SK I-5.
15. $(\mathrm{SO}_n \times \mathrm{GL}_m, \omega_1 \otimes \omega_1, \mathbb{C}^n \otimes \mathbb{C}^m)$ for $n \geq 3, \frac{1}{2}n \geq m \geq 1$.
We have $G_v^\circ \cong \mathrm{SO}_m \times \mathrm{SO}_{n-m}$ and $f(X) = \det(X^\top QX)$ for $X \in \mathrm{Mat}_{n, m} \cong \mathbb{C}^n \otimes \mathbb{C}^m$, $\deg(f) = 2m$, where $Q = g^\top Qg$ for $g \in \mathrm{SO}_n$.
16. $(\mathrm{GL}_1 \times \mathrm{Spin}_7, \mu \otimes \text{spinrep}, \mathbb{C} \otimes V^8)$.
We have $G_v^\circ \cong \mathrm{G}_2$ and $\deg(f) = 2$, where f is the relative invariant of SK I-15 for $m = 1, n = 8$.
17. $(\mathrm{GL}_2 \times \mathrm{Spin}_7, \omega_1 \otimes \text{spinrep}, \mathbb{C}^2 \otimes V^8)$.
We have $G_v^\circ \cong \mathrm{SO}_2 \times \mathrm{SL}_3$ and $\deg(f) = 4$, where f is the relative invariant of SK I-15 for $m = 2, n = 8$.
18. $(\mathrm{GL}_3 \times \mathrm{Spin}_7, \omega_1 \otimes \text{spinrep}, \mathbb{C}^3 \otimes V^8)$.
We have $G_v^\circ \cong \mathrm{SO}_3 \times \mathrm{SL}_2$ and $\deg(f) = 6$, where f is the relative invariant of SK I-15 for $m = 3, n = 8$.
19. $(\mathrm{GL}_1 \times \mathrm{Spin}_9, \mu \otimes \text{spinrep}, \mathbb{C} \otimes V^{16})$.
We have $G_v^\circ \cong \mathrm{Spin}_7$ and $\deg(f) = 2$.
20. $(\mathrm{GL}_2 \times \mathrm{Spin}_{10}, \omega_1 \otimes \text{halfspinrep}, \mathbb{C}^2 \otimes V^{16})$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{G}_2$ and $\deg(f) = 4$.
21. $(\mathrm{GL}_3 \times \mathrm{Spin}_{10}, \omega_1 \otimes \text{halfspinrep}, \mathbb{C}^3 \otimes V^{16})$.
We have $G_v^\circ \cong \mathrm{SO}_3 \times \mathrm{SL}_2$ and $\deg(f) = 12$.
22. $(\mathrm{GL}_1 \times \mathrm{Spin}_{11}, \mu \otimes \text{spinrep}, \mathbb{C} \otimes V^{32})$.
We have $G_v^\circ \cong \mathrm{SL}_5$ and $\deg(f) = 4$.
23. $(\mathrm{GL}_1 \times \mathrm{Spin}_{12}, \mu \otimes \text{halfspinrep}, \mathbb{C} \otimes V^{32})$.
We have $G_v^\circ \cong \mathrm{SL}_6$ and $\deg(f) = 4$.
24. $(\mathrm{GL}_1 \times \mathrm{Spin}_{14}, \mu \otimes \text{halfspinrep}, \mathbb{C} \otimes V^{64})$.
We have $G_v^\circ \cong \mathrm{G}_2 \times \mathrm{G}_2$ and $\deg(f) = 8$.
25. $(\mathrm{GL}_1 \times \mathrm{G}_2, \mu \otimes \omega_2, \mathbb{C} \otimes V^7)$.
We have $G_v^\circ \cong \mathrm{SL}_3$ and $\deg(f) = 2$, where f is the relative invariant of SK I-15 for $m = 1, n = 7$.
26. $(\mathrm{GL}_2 \times \mathrm{G}_2, \omega_1 \otimes \omega_2, \mathbb{C}^2 \otimes V^7)$.
We have $G_v^\circ \cong \mathrm{GL}_2$ and $\deg(f) = 4$, where f is the relative invariant of SK I-15 for $m = 2, n = 7$.
27. $(\mathrm{GL}_1 \times \mathrm{E}_6, \mu \otimes \omega_1, \mathbb{C} \otimes V^{27})$.
We have $G_v^\circ \cong \mathrm{F}_4$ and $\deg(f) = 4$.
28. $(\mathrm{GL}_2 \times \mathrm{E}_6, \omega_1 \otimes \omega_1, \mathbb{C}^2 \otimes V^{27})$.
We have $G_v^\circ \cong \mathrm{SO}_8$ and $\deg(f) = 12$.
29. $(\mathrm{GL}_1 \times \mathrm{E}_7, \mu \otimes \omega_6, \mathbb{C} \otimes V^{56})$.
We have $G_v^\circ \cong \mathrm{E}_6$ and $\deg(f) = 4$.

SK II Non-regular irreducible reduced prehomogeneous modules with non-constant relative invariant.

1. $(\mathrm{GL}_1 \times \mathrm{Sp}_n \times \mathrm{SO}_3, \mu \otimes \omega_1 \otimes \omega_1, \mathbb{C} \otimes \mathbb{C}^{2n} \otimes \mathbb{C}^3)$.
We have $G_v^\circ \cong (\mathrm{Sp}_{n-2} \times \mathrm{SO}_2) \cdot \mathrm{Un}_{2n-3}$ and $f(X) = \mathrm{trace}(X^\top JXQ)^2$ for $X \in \mathrm{Mat}_{2n,3} \cong \mathbb{C} \otimes \mathbb{C}^{2n} \otimes \mathbb{C}^3$.

SK III Non-regular irreducible reduced prehomogeneous modules without non-constant relative invariants.

1. $(G \times \mathrm{GL}_m, \varrho \otimes \omega_1, V^n \otimes \mathbb{C}^m)$,
where $\varrho : G \rightarrow \mathrm{GL}(V^n)$ is an n -dimensional irreducible representation of a semisimple algebraic group $G (\neq \mathrm{SL}_n)$ with $m > n \geq 3$. We have $G_v^\circ \cong (G \times \mathrm{GL}_{m-n}) \cdot G_{n(m-n)}^+$. The module $(G \times \mathrm{SL}_m, \varrho \otimes \omega_1)$ is prehomogeneous with $G_v^\circ \cong (G \times \mathrm{SL}_{m-n}) \cdot G_{n(m-n)}^+$.
2. $(\mathrm{SL}_n \times \mathrm{GL}_m, \omega_1 \otimes \omega_1, \mathbb{C}^n \otimes \mathbb{C}^m)$ for $\frac{1}{2}m \geq n \geq 1$.
We have $G_v^\circ \cong (\mathrm{SL}_n \times \mathrm{GL}_{m-n}) \cdot G_{n(m-n)}^+$. The module $(\mathrm{SL}_n \times \mathrm{SL}_m, \omega_1 \otimes \omega_1)$ is prehomogeneous with $G_v^\circ \cong (\mathrm{SL}_n \times \mathrm{SL}_{m-n}) \cdot G_{n(m-n)}^+$.
3. $(\mathrm{GL}_{2n+1}, \omega_2, \bigwedge^2 \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{Sp}_n \times \mathrm{GL}_1) \cdot G_{2n}^+$. The module $(\mathrm{SL}_{2n+1}, \omega_2)$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_n \cdot G_{2n}^+$.
4. $(\mathrm{GL}_2 \times \mathrm{SL}_{2n+1}, \omega_1 \otimes \omega_2, \mathbb{C}^2 \otimes \bigwedge^2 \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SL}_2) \cdot G_{2n}^+$ (see lemma 1.4 in Kimura et al. [4]). The module $(\mathrm{SL}_2 \times \mathrm{SL}_{2n+1}, \omega_1 \otimes \omega_2)$ is prehomogeneous with $G_v^\circ \cong \mathrm{SL}_2 \cdot G_{2n}^+$.
5. $(\mathrm{Sp}_n \times \mathrm{GL}_{2m+1}, \omega_1 \otimes \omega_1, \mathbb{C}^{2n} \otimes \mathbb{C}^{2m+1})$ for $n > 2m + 1 \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_m \times \mathrm{Sp}_{n-m}) \cdot \mathrm{Un}_{2n-1}$. The module $(\mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, \omega_1 \otimes \omega_1)$ is prehomogeneous with $G_v^\circ \cong (\mathrm{Sp}_m \times \mathrm{Sp}_{n-m}) \cdot \mathrm{Un}_{2n-1}$.
6. $(\mathrm{GL}_1 \times \mathrm{Spin}_{10}, \mu \otimes \text{halfspinrep}, \mathbb{C} \otimes V^{16})$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Spin}_7) \cdot G_8^+$. The module $(\mathrm{Spin}_{10}, \text{halfspinrep})$ is prehomogeneous with $G_v^\circ \cong \mathrm{Spin}_7 \cdot G_8^+$.

1.2 Non-irreducible simple prehomogeneous modules

The simple prehomogeneous, including the non-irreducible ones, were classified by Kimura, thus we will label them by Ks n , where n is the number of the module in §3 of Kimura's article [3].

In this section, it is understood that each representation ϱ_i of the simple group is composed with a scalar multiplication μ of GL_1^k . We shall simply write ϱ_i instead of $\mu \otimes \varrho_i$. In some cases, a module $V_1 \oplus \dots \oplus V_k$ will be prehomogeneous even with fewer than k scalar multiplications, in which case we will state this fact explicitly. We shall also state the connected component G_v° of the generic isotropy subgroup and the relative invariants $f_1, \dots, f_l$ where they exist.

Let $G = \mathrm{GL}_1^k \times G_s$ be a reductive group, where G_s is a simple algebraic group, let $(\varrho_1, V_1), \dots, (\varrho_k, V_k)$ be irreducible G_s -modules and $V = V_1 \oplus \dots \oplus V_k$ a G_s -module with representation $\varrho = \varrho_1 \oplus \dots \oplus \varrho_k$. Then (G, ϱ, V) is equivalent to one of the following:

Ks I Regular non-irreducible simple prehomogeneous modules.

1. $(\mathrm{GL}_1^2 \times \mathrm{SL}_n, \omega_1 \oplus \omega_1^*, \mathbb{C}^n \oplus \mathbb{C}^{n*})$ for $n \geq 3$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{SL}_{n-1}$ and $f_1(x, y) = \langle x|y \rangle$, where $(x, y) \in \mathbb{C}^n \oplus \mathbb{C}^{n*}$ and $\langle \cdot | \cdot \rangle$ is the dual pairing. The module $(\mathrm{GL}_1 \times \mathrm{SL}_n, (\mu \otimes \omega_1) \oplus \omega_1^*)$ is prehomogeneous with $G_v^\circ = \mathrm{SL}_{n-1}$.

2. $(\mathrm{GL}_1^n \times \mathrm{SL}_n, \omega_1^{\oplus n}, (\mathbb{C}^n)^{\oplus n})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1^{n-1}$ and $f_1(X) = \det(X)$ for $X \in \mathrm{Mat}_n \cong (\mathbb{C}^n)^{\oplus n}$. The module $(\mathrm{GL}_1 \times \mathrm{SL}_n, \mu \otimes \omega_1^{\oplus n})$ is prehomogeneous with $G_v^\circ = \{1\}$.
3. $(\mathrm{GL}_1^{n+1} \times \mathrm{SL}_n, \omega_1^{\oplus n+1}, (\mathbb{C}^n)^{\oplus n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong \{1\}$ and $f_i(X) = \det(x_1, \dots, \cancel{x_i}, \dots, x_{n+1})$ for $X = (x_1, \dots, x_{n+1}) \in \mathrm{Mat}_{n,n+1} \cong (\mathbb{C}^n)^{\oplus n+1}$.
4. $(\mathrm{GL}_1^{n+1} \times \mathrm{SL}_n, \omega_1^{\oplus n} \oplus \omega_1^*, (\mathbb{C}^n)^{\oplus n} \oplus \mathbb{C}^{n*})$ for $n \geq 3$.
We have $G_v^\circ \cong \{1\}$ and $f_1(X) = \langle x_1 | y \rangle, \dots, f_n(X) = \langle x_n | y \rangle, f_{n+1}(X) = \det(x_1, \dots, x_n)$ for $X = (x_1, \dots, x_n, y) \in (\mathbb{C}^n)^{\oplus n} \oplus \mathbb{C}^{n*}$.
5. $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1 \oplus \omega_1, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_{n-1}$ and $f_1(X, y, z) = \mathrm{Pf}(X), f_2(X, y, z) = y^\top X^\# z$, where $(X, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n}$ and $X^\#$ is the cofactor matrix of X . The module $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n}, (\mu \otimes \omega_2) \oplus (\mu \otimes (\omega_1 \oplus \omega_1)))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1}$.
6. $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1 \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_{n-1}$ and $f_1(X, y, z) = \mathrm{Pf}(X), f_2(X, y, z) = \langle y | z \rangle$, where $(X, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*}$. The module $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n}, (\mu \otimes \omega_2) \oplus (\mu \otimes (\omega_1 \oplus \omega_1^*)))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1}$.
7. $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1^* \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*})$ for $n \geq 3$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_{n-1}$ and $f_1(X, y, z) = \mathrm{Pf}(X), f_2(X, y, z) = y^\top X z$, where $(X, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*}$. The module $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n}, (\mu \otimes \omega_2) \oplus (\mu \otimes (\omega_1^* \oplus \omega_1^*)))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1}$.
8. $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_n$. The module $(\mathrm{GL}_1 \times \mathrm{Sp}_{2n+1}, \mu \otimes (\omega_2 \oplus \omega_1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_n$, see p. 94 in [3] for the relative invariant.
9. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1 \oplus \omega_1 \oplus \omega_1, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-1}$, see p. 94 in [3] for the relative invariants.
10. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1 \oplus \omega_1^* \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1*} \oplus \mathbb{C}^{2n+1*})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-1}$ and $f_1(X) = \mathrm{Pf} \begin{pmatrix} A & x \\ x^\top & 0 \end{pmatrix}, f_2(X) = \langle x | y \rangle, f_3(X) = \langle x | z \rangle, f_4 = y^\top A z$ for $X = (A, x, y, z) \in \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1*} \oplus \mathbb{C}^{2n+1*}$.
11. $(\mathrm{GL}_1^2 \times \mathrm{SL}_n, 2\omega_1 \oplus \omega_1, \mathrm{Sym}^2 \mathbb{C}^n \oplus \mathbb{C}^n)$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{SO}_{n-1}$ and $f_1(X) = \det(A), f_2(X) = x^\top A^\# x$ for $X = (A, x) \in \mathrm{Sym}^2 \mathbb{C}^n \oplus \mathbb{C}^n$.
12. $(\mathrm{GL}_1^2 \times \mathrm{SL}_n, 2\omega_1 \oplus \omega_1^*, \mathrm{Sym}^2 \mathbb{C}^n \oplus \mathbb{C}^{n*})$ for $n \geq 3$.
We have $G_v^\circ \cong \mathrm{SO}_{n-1}$ and $f_1(X) = \det(A), f_2(X) = x^\top A x$ for $X = (A, x) \in \mathrm{Sym}^2 \mathbb{C}^n \oplus \mathbb{C}^n$.
13. $(\mathrm{GL}_1^2 \times \mathrm{SL}_7, \omega_3 \oplus \omega_1, \bigwedge^3 \mathbb{C}^7 \oplus \mathbb{C}^7)$.
We have $G_v^\circ \cong \mathrm{SL}_3$, see p. 96 in [3] for the relative invariants.
14. $(\mathrm{GL}_1^2 \times \mathrm{SL}_7, \omega_3 \oplus \omega_1^*, \bigwedge^3 \mathbb{C}^7 \oplus \mathbb{C}^{7*})$.
We have $G_v^\circ \cong \mathrm{SL}_3$, see p. 96 in [3] for the relative invariants.
15. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_8, \mathrm{spinrep} \oplus \mathrm{halfspinrep}, V^8 \oplus V^8)$.
We have $G_v^\circ \cong \mathrm{G}_2$ and two quadratic invariants $f_1(x, y) = q_1(x), f_2(x, y) = q_2(y)$ for $(x, y) \in V^8 \oplus V^8$.

16. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_7, \mathrm{vecrep} \oplus \mathrm{spinrep}, V^7 \oplus V^8)$.
We have $G_v^\circ \cong \mathrm{SL}_3$ and two quadratic invariants $f_1(x, y) = q_1(x)$, $f_2(x, y) = q_2(y)$ for $(x, y) \in V^7 \oplus V^8$.
17. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10}, \mathrm{halfspinrep}_{\mathrm{even}} \oplus \mathrm{halfspinrep}_{\mathrm{even}}, V^{16} \oplus V^{16})$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{G}_2$, see p. 96 in [3] for the relative invariants. The module $(\mathrm{GL}_1 \times \mathrm{Spin}_{10}, \mu \otimes (\mathrm{halfspinrep}_{\mathrm{even}} \oplus \mathrm{halfspinrep}_{\mathrm{even}}))$ is prehomogeneous with $G_v^\circ \cong \mathrm{G}_2$.
18. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10}, \mathrm{vecrep} \oplus \mathrm{halfspinrep}, V^{10} \oplus V^{16})$.
We have $G_v^\circ \cong \mathrm{Spin}_7$, see p. 97 in [3] for the relative invariants.
19. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{12}, \mathrm{vecrep} \oplus \mathrm{halfspinrep}, V^{12} \oplus V^{32})$.
We have $G_v^\circ \cong \mathrm{SL}_5$, see p. 97 in [3] for the relative invariants.
20. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n, \omega_1 \oplus \omega_1, \mathbb{C}^{2n} \oplus \mathbb{C}^{2n})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_{n-1}$, see p. 97 in [3] for the relative invariant. The module $(\mathrm{GL}_1 \times \mathrm{Sp}_n, \mu \otimes (\omega_1 \oplus \omega_1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1}$.
21. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_3, \omega_3 \oplus \omega_1, V^{14} \oplus \mathbb{C}^6)$.
We have $G_v^\circ \cong \mathrm{SL}_2$, see p. 97 in [3] for the relative invariants.

Ks II Non-regular non-irreducible simple prehomogeneous modules.

1. $(\mathrm{GL}_1^k \times \mathrm{SL}_n, \omega_1^{\oplus k}, (\mathbb{C}^n)^{\oplus k})$ for $2 \leq k \leq n-1$.
We have $G_v^\circ \cong (\mathrm{GL}_1^k \times \mathrm{SL}_{n-k}) \cdot \mathrm{G}_{k(n-k)}^+$. The module $(\mathrm{SL}_n, \omega_1^{\oplus k})$ is prehomogeneous with $G_v^\circ \cong \mathrm{SL}_{n-k} \cdot \mathrm{G}_{k(n-k)}^+$.
2. $(\mathrm{GL}_1^k \times \mathrm{SL}_n, \omega_1^{\oplus k-1} \oplus \omega_1^*, (\mathbb{C}^n)^{\oplus k-1} \oplus \mathbb{C}^{n*})$ for $3 \leq k \leq n$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SL}_{n-k+1}) \cdot \mathrm{G}_{(n-k+1)(k-2)}^+$ and $f_1(X) = \langle x_1 | y \rangle, \dots, f_{k-1}(X) = \langle x_{k-1} | y \rangle$ for $X = (x_1, \dots, x_{k-1}, y) \in (\mathbb{C}^n)^{\oplus k-1} \oplus \mathbb{C}^{n*}$. The module $(\mathrm{GL}_1^{k-1} \times \mathrm{SL}_n, (\mu \otimes \omega_1^{\oplus k-1}) \oplus \omega_1^*)$ is prehomogeneous with $G_v^\circ \cong \mathrm{SL}_{n-k+1} \cdot \mathrm{G}_{(n-k+1)(k-2)}^+$.
3. $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_2, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \bigwedge^2 \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1^2 \cdot \mathrm{G}_{2n}^+$. The module $(\mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_2)$ is prehomogeneous with $G_v^\circ \cong \mathrm{G}_{2n}^+$.
4. $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-1}) \cdot \mathrm{Un}_{n-1}$ and $f_1(X) = \mathrm{Pf}(X)$ where $X \in \bigwedge^2 \mathbb{C}^{2n}$. The module $(\mathrm{GL}_1 \times \mathrm{SL}_{2n}, \mu \otimes (\omega_2 \oplus \omega_1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1} \cdot \mathrm{Un}_{n-1}$.
5. $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*})$ for $n \geq 3$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-1}) \cdot \mathrm{Un}_{n-1}$ and $f_1(X) = \mathrm{Pf}(X)$ where $X \in \bigwedge^2 \mathbb{C}^{2n}$. The module $(\mathrm{GL}_1 \times \mathrm{SL}_{2n}, \mu \otimes (\omega_2 \oplus \omega_1^*))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1} \cdot \mathrm{Un}_{n-1}$.
6. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1 \oplus \omega_1 \oplus \omega_1, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$ and $f_1(X) = \mathrm{Pf}(A)$, $f_2(X) = x^\top A^\# y$, $f_3(X) = y^\top A^\# z$, $f_4(X) = z^\top A x$ for $X = (A, x, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n}$.
7. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1 \oplus \omega_1 \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$ and $f_1(X) = \mathrm{Pf}(A)$, $f_2(X) = x^\top A^\# y$, $f_3(X) = \langle x | z \rangle$, $f_4(X) = \langle y | z \rangle$ for $X = (A, x, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*}$.
8. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1 \oplus \omega_1^* \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*})$ for $n \geq 3$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$ and $f_1(X) = \mathrm{Pf}(A)$, $f_2(X) = \langle x | y \rangle$, $f_3(X) = \langle x | z \rangle$, $f_4(X) = y^\top A z$ for $X = (A, x, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*}$.

9. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n}, \omega_2 \oplus \omega_1^* \oplus \omega_1^* \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*})$ for $n \geq 3$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$ and $f_1(X) = \mathrm{Pf}(A)$, $f_2(X) = x^\top Ay$, $f_3(X) = y^\top Az$, $f_4(X) = z^\top Ax$ for $X = (A, x, y, z) \in \bigwedge^2 \mathbb{C}^{2n} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*} \oplus \mathbb{C}^{2n*}$.
10. $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1*})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1^2 \times \mathrm{Sp}_{n-1}) \cdot \mathrm{Un}_{4n-2}$. The module $(\mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1^*)$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1} \cdot \mathrm{Un}_{4n-2}$.
11. $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1 \oplus \omega_1, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-1}) \cdot \mathrm{Un}_{2n-1}$, see p. 99 in [3] for the relative invariants. The module $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1}, (\mu \otimes (\omega_2 \oplus \omega_1)) \oplus (\mu \otimes \omega_1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1} \cdot \mathrm{Un}_{2n-1}$.
12. $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1 \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1*})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-1}) \cdot \mathrm{Un}_{2n-1}$, see p. 99 in [3] for the relative invariants. The module $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1}, (\mu \otimes (\omega_2 \oplus \omega_1)) \oplus (\mu \otimes \omega_1^*))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1} \cdot \mathrm{Un}_{2n-1}$.
13. $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1^* \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1*} \oplus \mathbb{C}^{2n+1*})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-1}) \cdot \mathrm{Un}_{2n-1}$ and $f_1(X) = x^\top Ay$ for $X = (A, x, y) \in \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1*} \oplus \mathbb{C}^{2n+1*}$. The module $(\mathrm{GL}_1 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus (\mu \otimes (\omega_1^* \oplus \omega_1^*)))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-1} \cdot \mathrm{Un}_{2n-2}$.
14. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus \omega_1^* \oplus \omega_1^* \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1} \oplus \mathbb{C}^{2n+1})$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-2}) \cdot \mathrm{Un}_{4n-6}$ and $f_1(X) = x^\top Ay$, $f_2(X) = y^\top Az$, $f_3(X) = z^\top Ax$. The module $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n+1}, \omega_2 \oplus (\mu \otimes \omega_1^*) \oplus (\mu \otimes \omega_1^*) \oplus (\mu \otimes \omega_1^*))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{4n-6}$.
15. $(\mathrm{GL}_1^2 \times \mathrm{SL}_6, \omega_3 \oplus \omega_1, \bigwedge^3 \mathbb{C}^6 \oplus \mathbb{C}^6)$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SL}_2 \times \mathrm{SL}_2) \cdot \mathrm{G}_4^+$, see p. 100 in [3] for the relative invariant. The module $(\mathrm{GL}_1 \times \mathrm{SL}_6, \mu \otimes (\omega_3 \oplus \omega_1))$ is prehomogeneous with $G_v^\circ \cong (\mathrm{SL}_2 \times \mathrm{SL}_2) \cdot \mathrm{G}_4^+$.
16. $(\mathrm{GL}_1^3 \times \mathrm{SL}_6, \omega_3 \oplus \omega_1 \oplus \omega_1, \bigwedge^3 \mathbb{C}^6 \oplus \mathbb{C}^6 \oplus \mathbb{C}^6)$.
We have $G_v^\circ \cong \mathrm{GL}_1^2 \cdot \mathrm{G}_4^+$, see p. 100 in [3] for the relative invariant. The module $(\mathrm{GL}_1 \times \mathrm{SL}_6, \mu \otimes (\omega_3 \oplus \omega_1 \oplus \omega_1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{G}_4^+$.
17. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n, \omega_1 \oplus \omega_1 \oplus \omega_1, \mathbb{C}^{2n} \oplus \mathbb{C}^{2n} \oplus \mathbb{C}^{2n})$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$, see p. 100 in [3] for the relative invariants.
18. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_2, \omega_2 \oplus \omega_1, V^5 \oplus \mathbb{C}^4)$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$, see p. 100 in [3] for the relative invariant. The module $(\mathrm{GL}_1 \times \mathrm{Sp}_2, (\mu \otimes \omega_2) \oplus \omega_1)$ is prehomogeneous with $G_v^\circ \cong \mathrm{Un}_2$.
19. $(\mathrm{GL}_1^3 \times \mathrm{SL}_5, \omega_2 \oplus \omega_2 \oplus \omega_1^*, \bigwedge^2 \mathbb{C}^5 \oplus \bigwedge^2 \mathbb{C}^5 \oplus \mathbb{C}^5)$.
See proposition 1.1 in [4].

1.3 Two-simple prehomogeneous modules of type I

In this and the following chapter we shall give a classification of the non-irreducible two-simple prehomogeneous modules, i.e. modules of the form

$$\begin{aligned}
 &(\mathrm{GL}_1^l \times G_1 \times G_2, \\
 &(\varrho_1 \otimes \tilde{\varrho}_1) \oplus \dots \oplus (\varrho_k \otimes \tilde{\varrho}_k) \oplus (\sigma_1 \otimes 1) \oplus \dots \oplus (\sigma_s \otimes 1) \oplus (1 \otimes \tau_1) \oplus \dots \oplus (1 \otimes \tau_t), \\
 &V_1 \oplus \dots \oplus V_l),
 \end{aligned}$$

where G_1 and G_2 are simple algebraic groups, $l = k + s + t$, and the ϱ_i , σ_j (resp. $\tilde{\varrho}_i$, τ_j) are irreducible representations of G_1 (resp. G_2). As in the previous chapter, it is understood that each of these representations is composed with a scalar multiplication of GL_1^k . First, we give the classification of the type I-modules, i.e. at least one of the modules $(\mathrm{GL}_1 \times G_1 \times G_2, \varrho_i \otimes \tilde{\varrho}_i)$ is a non-trivial prehomogeneous module. These were classified by Kimura et al. [4], thus we shall refer to them as KI n , where n is the number of the module in §3 of [4]. We shall state the non-irreducible modules only, as the irreducible ones already appear in the table SK or as castling transformas of those (see also theorem 1.5 in [4]). In the next chapter, we shall classify the remaining two-simple modules of type II.

Let (G, ϱ, V) be a two-simple prehomogeneous module of type I. Then it is equivalent to one of the following:

KI I Regular two-simple prehomogeneous modules of type I.

1. $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes \omega_1))$.
We have $G_v^\circ \cong \{1\}$.
2. $(\mathrm{GL}_1^3 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_1$. The module $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1))$ is prehomogeneous with $G_v^\circ \cong \{1\}$.
3. $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_3, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong \mathrm{SO}_3$.
4. $(\mathrm{GL}_1^3 \times \mathrm{SL}_4 \times \mathrm{SL}_3, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{SO}_2$.
5. $(\mathrm{GL}_1^3 \times \mathrm{SL}_4 \times \mathrm{SL}_4, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SO}_2$.
6. $(\mathrm{GL}_1^3 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (\omega_1^{(*)} \otimes 1))$.
We have $G_v^\circ \cong \{1\}$.
7. $(\mathrm{GL}_1^2 \times \mathrm{SL}_5 \times \mathrm{SL}_3, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{SO}_2$.
8. $(\mathrm{GL}_1^2 \times \mathrm{SL}_5 \times \mathrm{SL}_8, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SO}_2$.
9. $(\mathrm{GL}_1^2 \times \mathrm{SL}_5 \times \mathrm{SL}_9, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{SL}_2 \times \mathrm{SL}_2$. The module $(\mathrm{GL}_1 \times \mathrm{SL}_5 \times \mathrm{SL}_9, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$ is prehomogeneous with $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SL}_2$.
10. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_{n-m} \times \mathrm{Sp}_{m-1}$. The module $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{Sp}_{2m}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)}))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_{n-m} \times \mathrm{Sp}_{m-1}$.
11. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_2, (\omega_1 \otimes \omega_1) \oplus (1 \otimes 2\omega_1))$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-1} \times \mathrm{SO}_2$.
12. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes 3\omega_1))$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-1}$.
13. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_2, (\omega_1 \otimes \omega_1) \oplus (1 \otimes 2\omega_2) \oplus (1 \otimes \omega_1))$. We have $G_v^\circ \cong \mathrm{Sp}_{n-1}$.
14. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{Sp}_m \times \mathrm{Sp}_{n-m-1}$. The module $(\mathrm{GL}_1 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{Sp}_m \times \mathrm{Sp}_{n-m-1}$.

15. $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes (\omega_1 \oplus \omega_1)^{(*)}))$.
We have $G_v^\circ \cong \mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-1}$.
16. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{GL}_1$. The module $(\mathrm{GL}_1^2 \times \mathrm{Sp}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1^*))$ is prehomogeneous with $G_v^\circ \cong \{1\}$.
17. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_2 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong \mathrm{SO}_2$.
18. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \{1\}$.
19. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_4, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \{1\}$.
20. $(\mathrm{GL}_1^2 \times \mathrm{SO}_n \times \mathrm{SL}_m, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{SO}_{m-1} \times \mathrm{SO}_{n-m}$.
21. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_7 \times \mathrm{SL}_2, (\mathrm{spinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{SL}_3$.
22. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_7 \times \mathrm{SL}_3, (\mathrm{spinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_2$.
23. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_7 \times \mathrm{SL}_6, (\mathrm{spinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_2$.
24. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_7 \times \mathrm{SL}_7, (\mathrm{spinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_3$.
25. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_7 \times \mathrm{SL}_2, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{spinrep} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_2$.
26. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_7 \times \mathrm{SL}_2, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{spinrep} \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{SL}_2$.
27. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_7 \times \mathrm{SL}_6, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{spinrep} \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_2$.
28. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_8 \times \mathrm{SL}_2, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{halfspinrep} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{SL}_3 \times \mathrm{SO}_2$.
29. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_8 \times \mathrm{SL}_3, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{halfspinrep} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_3$.
30. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_8 \times \mathrm{SL}_2, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{halfspinrep} \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{SL}_3$.
31. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_8 \times \mathrm{SL}_3, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{halfspinrep} \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_2$.
32. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_8 \times \mathrm{SL}_6, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{halfspinrep} \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_2$.
33. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_8 \times \mathrm{SL}_7, (\mathrm{vecrep} \otimes \omega_1) \oplus (\mathrm{halfspinrep} \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_3$.
34. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\mathrm{halfspinrep} \otimes \omega_1) \oplus (1 \otimes 2\omega_1))$.
We have $G_v^\circ \cong \mathrm{G}_2 \times \mathrm{SO}_3$.
35. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\mathrm{halfspinrep} \otimes \omega_1) \oplus (1 \otimes 3\omega_1))$.
We have $G_v^\circ \cong \mathrm{G}_2$.

36. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{G}_2$. The module $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ is prehomogeneous with $G_v^\circ \cong \mathrm{G}_2$.
37. $(\mathrm{GL}_1^3 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes 2\omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{G}_2$.
38. $(\mathrm{GL}_1^4 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{G}_2$.
39. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_3, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_2$.
40. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_{14}, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SO}_2$.
41. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_{15}, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \times \mathrm{SL}_4$. The module $(\mathrm{GL}_1 \times \mathrm{Spin}_{10} \times \mathrm{SL}_{15}, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}))$ is prehomogeneous with $G_v^\circ \cong \mathrm{SL}_4$.
42. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\text{vecrep} \otimes \omega_1) \oplus (\text{halfspinrep} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{G}_2$.
43. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_3, (\text{halfspinrep} \otimes \omega_1) \oplus (\text{vecrep} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{SL}_3 \times \mathrm{SO}_2$.
44. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_4, (\text{halfspinrep} \otimes \omega_1) \oplus (\text{vecrep} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{SL}_2 \times \mathrm{SL}_2$.
45. $(\mathrm{GL}_1^2 \times \mathrm{G}_2 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{SL}_2$.
46. $(\mathrm{GL}_1^2 \times \mathrm{G}_2 \times \mathrm{SL}_6, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{SL}_2$.

KI II Non-regular two-simple prehomogeneous modules of type I.

1. (a) $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1} \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1^2 \cdot \mathrm{G}_1^+$.
(b) $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1} \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes 2\omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SO}_2) \cdot \mathrm{G}_{2n}^+$.
(c) $(\mathrm{GL}_1^2 \times \mathrm{SL}_{2n+1} \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes 3\omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{G}_n^+$.
2. (a) $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n+1} \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1^2 \cdot \mathrm{G}_{2n}^+$.
(b) $(\mathrm{GL}_1^3 \times \mathrm{SL}_{2n+1} \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes 2\omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{G}_{2n}^+$.
3. $(\mathrm{GL}_1^4 \times \mathrm{SL}_{2n+1} \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{G}_{2n}^+$.
4. $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SO}_2) \cdot \mathrm{Un}_2$.
5. $(\mathrm{GL}_1^3 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.
6. $(\mathrm{GL}_1^3 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.

7. (a) $(\mathrm{GL}_1^2 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SO}_2) \cdot \mathrm{Un}_2$.
- (b) $(\mathrm{GL}_1^2 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_1^2 \cdot \mathrm{Un}_2$.
8. (a) $(\mathrm{GL}_1^3 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.
- (b) $(\mathrm{GL}_1^3 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1^2 \cdot \mathrm{Un}_2$.
9. $(\mathrm{GL}_1^3 \times \mathrm{SL}_5 \times \mathrm{SL}_9, (\omega_2 \otimes \omega_1) \oplus (\omega_1^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^*))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.
10. $(\mathrm{GL}_1^3 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes 2\omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.
11. $(\mathrm{GL}_1^4 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.
12. $(\mathrm{GL}_1^2 \times \mathrm{SL}_6 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^{(*)} \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_3$.
13. (a) $(\mathrm{GL}_1^2 \times \mathrm{SL}_7 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_3$.
- (b) $(\mathrm{GL}_1^2 \times \mathrm{SL}_7 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1))$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SO}_2) \cdot \mathrm{Un}_2$.
14. $(\mathrm{GL}_1^3 \times \mathrm{SL}_7 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_2$.
15. $(\mathrm{GL}_1^2 \times \mathrm{SL}_9 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1))$.
We have $G_v^\circ \cong \mathrm{GL}_1 \cdot \mathrm{Un}_3$.
16. (a) $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2}$.
- (b) $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2}$.
17. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2m-2}$.
18. (a) $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)}))$ for $n > m \geq 2$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-2} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-4}$.
- (b) $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_2, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$.
- (c) $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)}))$ for $n > m \geq 2$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-2} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-4}$.
19. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_2, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes 2\omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$.
20. (a) $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_m \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2m-1}$.
- (b) $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1^2 \times \mathrm{Sp}_m \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2m-3}$.

- (c) $(\mathrm{GL}_1^2 \times \mathrm{Sp}_{m+1} \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$ for $m \geq 1$.
We have $G_v^\circ \cong (\mathrm{GL}_1^2 \times \mathrm{Sp}_{m-1}) \cdot \mathrm{Un}_{4m-1}$.
- (d) $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2))$ for $n > m + 1 \geq 2$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{SO}_2^m \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2m-1}$.
21. (a) $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1))$ for $n > m + 1$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-2}) \cdot \mathrm{Un}_{2n-4}$.
- (b) $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$ for $n > m + 1$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-2}) \cdot \mathrm{Un}_{2n-4}$.
- (c) $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)}))$ for $n > m + 1$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-2}) \cdot \mathrm{Un}_{2n-4}$.
22. (a) $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes (\omega_1 \oplus \omega_1)^{(*)}))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-1} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n-2m-1}$.
- (b) $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_{2m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*))$ for $n > m \geq 1$.
We have $G_v^\circ \cong (\mathrm{Sp}_{m-2} \times \mathrm{Sp}_{n-m-1}) \cdot \mathrm{Un}_{2n+2m-7}$.
- (c) $(\mathrm{GL}_1^4 \times \mathrm{Sp}_n \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$.
23. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes 2\omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong (\mathrm{SO}_2 \times \mathrm{Sp}_{n-2}) \cdot \mathrm{Un}_{2n-3}$.
24. $(\mathrm{GL}_1^3 \times \mathrm{Sp}_n \times \mathrm{SL}_5, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1^*))$ for $n \geq 3$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{Sp}_{n-3}) \cdot \mathrm{Un}_{2n-4}$.
25. $(\mathrm{GL}_1^2 \times \mathrm{Sp}_n \times \mathrm{SL}_2, (\omega_1 \otimes 2\omega_1) \oplus (1 \otimes \omega_1))$ for $n \geq 2$.
We have $G_v^\circ \cong \mathrm{Sp}_{n-2} \cdot \mathrm{Un}_{2n-3}$.
26. $(\mathrm{GL}_1^2 \times \mathrm{Spin}_{10} \times \mathrm{SL}_2, (\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
We have $G_v^\circ \cong (\mathrm{GL}_1 \times \mathrm{G}_2) \cdot \mathrm{G}_1^+$.

1.4 Two-simple prehomogeneous modules of type II

In this chapter we give a classification of the two-simple prehomogeneous modules of type II, i.e. modules of the form

$$(\mathrm{GL}_1^l \times G_1 \times G_2, \\ (\varrho_1 \otimes \tilde{\varrho}_1) \oplus \dots \oplus (\varrho_k \otimes \tilde{\varrho}_k) \oplus (\sigma_1 \otimes 1) \oplus \dots \oplus (\sigma_s \otimes 1) \oplus (1 \otimes \tau_1) \oplus \dots \oplus (1 \otimes \tau_t), \\ V_1 \oplus \dots \oplus V_l),$$

where all of the modules $(\mathrm{GL}_1 \times G_1 \times G_2, \varrho_i \otimes \tilde{\varrho}_i)$ are trivial prehomogeneous modules (see Kimura [2]). Note that we consider non-irreducible modules only. These were classified by Kimura et al. [5], thus we shall refer to them as KII n , where n is the number of the module in §5 of [5]. Unfortunately, it is not always obvious from the classification in which cases a module would be prehomogeneous even with fewer than l scalar multiplications.

Any indecomposable two-simple prehomogeneous module of type II is equivalent to one of the following:

KII I Two-simple prehomogeneous modules of type II obtained directly from any given simple module $(\mathrm{GL}_1^l \times G, \varrho_1 \oplus \dots \oplus \varrho_l)$.

1. For any representation $\sigma_1 \oplus \dots \oplus \sigma_s$ of G and $n \geq \sum_{i=1}^s \dim(\sigma_i)$:

$$(\mathrm{GL}_1^{l+s} \times G \times \mathrm{SL}_n, \\ (\sigma_1 \otimes \omega_1) \oplus \dots \oplus (\sigma_s \otimes \omega_1) \oplus (\varrho_1 \otimes 1) \oplus \dots \oplus (\varrho_l \otimes 1)).$$

2. For $t \geq 0$, $1 \leq k \leq l$ and $n = t - 1 + \sum_{i=1}^k \dim(\varrho_i)$:

$$(\mathrm{GL}_1^{l+t} \times G \times \mathrm{SL}_n, \\ (\varrho_1 \otimes \omega_1) \oplus \dots \oplus (\varrho_k \otimes \omega_1) \oplus (\varrho_{k+1}^* \otimes 1) \oplus \dots \oplus (\varrho_l^* \otimes 1) \oplus (1 \otimes \omega_1^{\oplus t})).$$

3. For $t \geq 1$, $1 \leq k \leq l$ and $n \geq t - 1 + \sum_{i=1}^k \dim(\varrho_i)$:

$$(\mathrm{GL}_1^{l+t} \times G \times \mathrm{SL}_n, \\ (\varrho_1 \otimes \omega_1) \oplus \dots \oplus (\varrho_k \otimes \omega_1) \oplus (\varrho_{k+1} \otimes 1) \oplus \dots \oplus (\varrho_l \otimes 1) \oplus (1 \otimes \omega_1^{\oplus t-1}) \oplus (1 \otimes \omega_1^*)).$$

KII II Two-simple prehomogeneous modules of type II of the form

$$(\mathrm{GL}_1^{k+s+t} \times G \times \mathrm{SL}_n, \\ (\varrho_1 \otimes \omega_1) \oplus \dots \oplus (\varrho_k \otimes \omega_1) \oplus (\sigma_1 \otimes 1) \oplus \dots \oplus (\sigma_s \otimes 1) \oplus (1 \otimes \tau_1) \oplus \dots \oplus (1 \otimes \tau_t)),$$

with $2 \leq \dim(\varrho_i) \leq n$ for all i and at least one $\tau_j \neq \omega_1^{(*)}$.

4. $G = \mathrm{SL}_m$ with $2 \leq m < n$.

4-i (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes 2\omega_1^{(*)})$.

(b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes 2\omega_1^{(*)}) \oplus (\omega_1^{(*)} \otimes 1)$.

(c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)})$.

(d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1^{(*)} \otimes 1)$.

(e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1^{(*)} \otimes 1) \oplus (\omega_1^{(*)} \otimes 1)$.

(f) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (1 \otimes \omega_1^{(*)})$.

(g) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^{(*)})$.

- 4-ii n even.

(a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus ((\omega_1 \oplus \omega_1)^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^{(*)})$.

(b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (\omega_1^* \otimes 1)$.

(c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus ((\omega_1 \oplus \omega_1 \oplus \omega_1)^{(*)} \otimes 1)$.

(d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1)$, m even.

(e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1^{(*)})$, m even.

(f) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_2 \otimes 1)$, m odd.

(g) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (1 \otimes \omega_1^{(*)}) \oplus (1 \otimes \omega_1^{(*)})$, m odd.

- 4-iii n odd.

(a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*)$, $m \geq 3$.

(b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_2^* \otimes 1)$.

(c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus ((\omega_1 \oplus \omega_1 \oplus \omega_1)^{(*)} \otimes 1)$, $m \geq 3$.

(d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus ((\omega_1 \oplus \omega_1)^{(*)} \otimes 1) \oplus (1 \otimes \omega_1)$.

(e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_2 \otimes 1)$, m even.

(f) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1)$, m even.

(g) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*)$, m even.

(h) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*)$, m even.

(i) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1)$, m odd.

(j) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1)$, m odd.

- (k) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1^*), m \text{ odd.}$
 - (l) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (\omega_1^* \otimes 1).$
 - (m) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus ((\omega_1 \oplus \omega_1 \oplus \omega_1)^{(*)} \otimes 1).$
 - (n) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus ((\omega_1 \oplus \omega_1)^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^{(*)}).$
 - (o) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1), m \text{ even.}$
 - (p) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*), m \text{ even.}$
 - (q) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1^{(*)}), m \text{ even.}$
 - (r) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (\omega_2 \otimes 1), m \text{ odd.}$
 - (s) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1), m \text{ odd.}$
 - (t) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*), m \text{ odd.}$
5. $G = \text{SL}_2, n > 2.$
- 5-i (a) $(2\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}).$
 - (b) $(2\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1 \otimes 1).$
 - 5-ii (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (2\omega_1 \otimes 1).$
 - (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (2\omega_1 \otimes 1) \oplus (1 \otimes \omega_1).$
 - (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (3\omega_1 \otimes 1), n \text{ even.}$
 - (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (2\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1), n \text{ even.}$
 - (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (2\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*), n \text{ even.}$
 - 5-iii (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*).$
 - (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (2\omega_1 \otimes 1).$
 - (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (3\omega_1 \otimes 1).$
 - (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (2\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1).$
 - (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (2\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^{(*)}).$
 - 5-iv $n = 5.$
 - (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_2^*).$
 - 5-v $n = 6.$
 - (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3).$
 - (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (1 \otimes \omega_1^*).$
 - (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1 \otimes 1).$
 - (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (2\omega_1 \otimes 1).$
 - (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (3\omega_1 \otimes 1).$
 - (f) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1).$
 - (g) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (2\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1).$
 - (h) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1).$
 - 5-vi $n = 7.$
 - (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3^{(*)}).$
 - (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3^{(*)}) \oplus (\omega_1 \otimes 1).$
6. $G = \text{SL}_3, n > 3.$
- (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (2\omega_1^{(*)} \otimes 1).$
 - (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_2), n = 5.$
7. $G = \text{SL}_4, n > 4.$
- 7-i $n \text{ odd.}$

- (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (2\omega_1 \otimes 1)$.
 (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_2 \otimes 1) \oplus (\omega_1^{(*)} \otimes 1)$.
 (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1^*)$.
- 7-ii $n = 5$.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_2)$.
- 7-iii $n = 6$.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3)$.
 (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1^* \otimes 1)$.
 (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_2^{(*)} \otimes 1)$.
 (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1^* \otimes 1) \oplus (\omega_1^* \otimes 1)$.
 (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (1 \otimes \omega_1)$.
8. $G = \mathrm{SL}_5$, $n > 5$.
 8-i n even.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_2 \otimes 1) \oplus (\omega_1^* \otimes 1)$.
 8-ii n odd.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (\omega_2 \otimes 1) \oplus (\omega_1^* \otimes 1)$.
 (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_2) \oplus (\omega_2^* \otimes 1) \oplus (\omega_1 \otimes 1)$.
 (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_2) \oplus (\omega_2^* \otimes 1) \oplus (1 \otimes \omega_1^*)$.
 8-iii $n = 6$.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3)$.
 (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1^{(*)} \otimes 1)$.
 (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_2^* \otimes 1)$.
 (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (\omega_1^* \otimes 1) \oplus (\omega_1^{(*)} \otimes 1)$.
 (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3) \oplus (1 \otimes \omega_1^*)$.
 8-iv $n = 7$.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3^{(*)})$.
9. $G = \mathrm{SL}_{2j}$, $n = 2j + 1$.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1)$.
 (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3^{(*)}) \oplus (\omega_1^{(*)} \otimes 1)$, $j = 3$ (i.e. $n = 7$).
10. $G = \mathrm{SL}_n$.

$$(\omega_1 \otimes \omega_1) \oplus (\varrho_1 \otimes 1) \oplus \dots \oplus (\varrho_k \otimes 1) \oplus (1 \otimes \varrho_{k+1}^*) \oplus \dots \oplus (1 \otimes \varrho_r^*),$$
 where $(\mathrm{GL}_1^r \times \mathrm{SL}_n, \varrho_1 \oplus \dots \oplus \varrho_r)$ is a simple prehomogeneous module.
11. $G = \mathrm{Sp}_m$, $2m < n$.
 11-i n odd.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2)$.
 (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_1 \otimes 1)$, $m = 2$.
 (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_2 \otimes 1)$, $m = 2$.
 (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1^*)$, $m = 2$.
 11-ii $n = 6$, $m = 2$.
 (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_3)$.

KII III Two-simple prehomogeneous modules of type II of the form

$$(\mathrm{GL}_1^{k+s+t} \times G \times \mathrm{SL}_n, \\ (\varrho_1 \otimes \omega_1) \oplus \dots \oplus (\varrho_k \otimes \omega_1) \oplus (\sigma_1 \otimes 1) \oplus \dots \oplus (\sigma_s \otimes 1) \oplus (1 \otimes \omega_1^{\oplus t})),$$

with $2 \leq \dim(\varrho_i) \leq n$ for all i and

$$(G, \varrho_1, \dots, \varrho_k, \sigma_1, \dots, \sigma_s) \neq (\mathrm{SL}_m, \omega_1, \dots, \omega_1, \omega_1^{(*)}, \dots, \omega_1^{(*)}).$$

12. $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_8, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes \omega_1)).$

13. $G = \mathrm{SL}_m.$

13-i $m < n.$

- (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^{(*)} \otimes 1).$
- (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^{(*)} \otimes 1) \oplus (\omega_1 \otimes 1).$
- (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^* \otimes 1) \oplus (\omega_1^* \otimes 1).$
- (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^*).$
- (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^* \otimes 1) \oplus (1 \otimes \omega_1).$
- (f) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2 \otimes 1) \oplus (\omega_1^* \otimes 1), m \text{ even.}$
- (g) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1), m \text{ even.}$
- (h) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_3 \otimes 1), m = 6.$
- (i) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_3 \otimes 1) \oplus (1 \otimes \omega_1), m = 6.$

13-ii $n = m + 1.$

- (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\omega_2 \otimes 1).$
- (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\omega_2^* \otimes 1), m \text{ even.}$
- (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\omega_3 \otimes 1), m = 6.$

13-iii $n \geq \frac{1}{2}m(m-1).$

- (a) $(\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*), m \text{ odd.}$
- (b) $(\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*), m \text{ odd, } n > \frac{1}{2}m(m-1).$
- (c) $(\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*), m = 5.$
- (d) $(\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1), m = 2j+1, n = 2j^2 + j + 1.$
- (e) $(\omega_2 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1), m = 2j, n = 2j^2 + j.$
- (f) $(\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1), m = 5, n = 10.$

14. $G = \mathrm{Sp}_m, n \geq 2m.$

- (a) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*).$
- (b) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_1 \otimes 1).$
- (c) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*).$
- (d) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1), n > 2m.$
- (e) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (\omega_1 \otimes 1), n = 2m.$
- (f) $(\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^{(*)}), n = 2m.$

15. $G = \mathrm{Spin}_{10}, n \geq 16.$

- (a) $(\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*).$
- (b) $(\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*), n \geq 17.$
- (c) $(\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1), n = 17.$
- (d) $(\text{halfspinrep} \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1), n = 16.$

KII IV Two-simple prehomogeneous modules of type II of the form

$$(\mathrm{GL}_1^{k+s_1+s_2+t_1+t_2} \times \mathrm{SL}_m \times \mathrm{SL}_n, \\ (\omega_1 \otimes 1)^{\oplus s_1} \oplus (\omega_1^* \otimes 1)^{\oplus s_2} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t_1} \oplus (1 \otimes \omega_1^*)^{\oplus t_2}),$$

where $n \geq m \geq 2$ and $k \geq 1$.

16. $n \geq km$.

16-i $n = m$. Then $k = 1$ and $1 \leq (s_1 + t_2) + (s_2 + t_1) \leq n + 1$, where one of $s_1 + t_2$ or $s_2 + t_1$ is 0 or 1.

16-ii $n = km, k \geq 2$.

(a) $t_1 = 0, 2 \leq t_2 \leq n, s_2 = 0, s_1 + kt_2 \leq m$.

(b) $t_2 = 0, 2 \leq t_1 \leq n, s_1 = 0, s_2 + kt_1 \leq m$.

16-iii $n = km + 1$. Then $t_1 \geq 3, t_2 = s_1 = 0, s_2 + k(t_1 - 1) \leq m$.

16-iv $n \geq km + t_1, n > km$.

(a) $k = 1, t_1 = 0, 2 \leq t_2 \leq n$ and $1 \leq (s_1 + t_2) + s_2 \leq m + 1$, where s_2 is 0 or 1.

(b) $k \geq 2, t_1 = 0, 2 \leq t_2 \leq n, s_2 = 0, s_1 + kt_2 \leq m$.

(c) $k \geq 1, t_1 = 1, 2 \leq t_2 \leq n, s_2 = 0, s_1 + kt_2 \leq m$.

17. $km > n$. These are the cases (17)-(25) in §5.4 of [5], but to keep things simple we subsume them under the case KII IV-17 here. See the following definition 1.1 for the definition of $T, \nu(k, m, n)$ and (a_i) . Also, we write $b_i = \frac{a_i}{a_{i+1}}$.

17-i (a) $t_2 \geq 1, s_2 = t_1 = 0, s_1 + kt_2 \leq m - b_j(n - t_2)$, where $(k, m, n) \in T$ and $j = \nu(k, m, n)$.

(b) $s_2 = t_2 = 0$ and let $p = km + t_1 - n (< m), q = kp - m (< n), (k, p, m) \in T$ (resp. $(k, q, p) \in T$) and $j = \nu(k, p, m)$ (resp. $j = (k, q, p)$).

i. $m \geq kp, s_1 = 0$ and $t_1 \leq p + 1$.

ii. $m \geq kp, s_1 = 1$ and $k + t_1 \leq p + 1$.

iii. $m \geq kp, 2 \leq s_1 \leq m$ and $t_1 + ks_1 \leq p$.

iv. $kp > m, s_1 \geq 1$ and $t_1 + ks_1 \leq p - b_j(m - s_1)$.

v. $kp > m, p \geq kq, s_1 = 0, t_1 = 1$ and $k \leq q + 1$.

vi. $kp > m, p \geq kq, s_1 = 0, 2 \leq t_1 \leq p$ and $kt_1 \leq q$.

vii. $kp > m, kq > p, s_1 = 0, t_1 \geq 1$ and $kt_1 \leq q - b_j(p - t_1)$.

(c) $t_2 = 0, s_2 \geq 1, s_1 = 0$ and let $p = km + t_1 - n (< m), q = kp + s_2 - m (< p), r = kq - p (< q), (k, q, p) \in T$ (resp. $(k, q, p) \in T, (k, r, q) \in T$) and $j = \nu(k, q, p)$ (resp. $j = \nu(k, r, q)$).

i. $m \geq kp + s_2$ and $t_1 \leq p + 1$.

ii. $m = kp + s_2 - 1, t_1 = 0, 1$ and $k + t_1 \leq p + 1$.

iii. $m = kp + 1, m \geq s_2 \geq 3, t_1 = 0$ and $k(s_2 - 1) \leq p$.

iv. $m = kp, m \geq s_2 \geq 2$ and $ks_2 \leq p$.

v. $kp > m, p \geq kq, t_1 = 0$ and $s_2 \leq q + 1$.

vi. $kp > m, p \geq kq, t_1 = 1$ and $s_2 + kt_1 \leq q + 1$.

vii. $kp > m, p \geq kq, p \geq t_1 \geq 2$ and $s_2 + kt_1 \leq q$.

viii. $kp > m, kq > p, t_1 \geq 1$ and $s_2 + kt_1 \leq q - b_j(p - t_1)$.

ix. $kp > m, kq > p, q \geq kr, t_1 = 0, s_2 = 1$ and $k \leq r + 1$.

- x. $kp > m, kq > p, q \geq kr, t_1 = 0, q \geq s_2 \geq 2$ and $ks_2 \leq r$.
- xi. $kp > m, kq > p, kr > q, t_1 = 0, s_2 \geq 1$ and $ks_2 \leq r - b_j(q - s_2)$.
- 17-ii (a) $t_2 = 1, s_2 = 0, t_1 \geq 1$ and let $p = km + t_1 - n - 1, (k, p, m) \in T$ and $j = \nu(k, p, m)$.
 - i. $m \geq kp$ and $(t_1 - 1) + k(k + s_1) \leq p$.
 - ii. $kp > m$ and $(t_1 - 1) + k(k + s_1) \leq p - b_j(m - k - s_1)$.
- (b) $t_2 = 0, s_2 \geq 1, s_1 = 1$ and let $p = km + t_1 - n, q = kp + s_2 - m - 1, (k, q, p) \in T$ and $j = \nu(k, q, p)$.
 - i. $kp > m, p \geq kq, (s_2 - 1) + k(k + t_1) \leq q$.
 - ii. $kp > m, kq > n, (s_2 - 1) + k(k + t_1) \leq q - b_j(p - k - t_1)$.
- 17-iii (a) $t_2 \geq 0, s_2 = 0, t_1 = 1, (s_1 + k) + k(t_2 - 1) \leq m - b_j(n - t_2)$ where $(k, m, n - 1) \in T$ and $j = \nu(k, m, n - 1)$.
- (b) $t_2 = 0, s_2 = 1, s_1 \geq 2$ and let $p = km + t_1 - n (< m), (k, p, m - 1) \in T$ and $j = \nu(k, p, m - 1)$.
 - i. $m \geq kp$ and $t_1 + ks_1 \leq p$.
 - ii. $kp > m$ and $t_1 + ks_1 \leq p - b_j(m - s_1)$.
- 17-iv (a) $t_2 = s_2 = 1$ and let $p = km + t_1 - n, (k, p, m - 1) \in T$ and $j = \nu(k, p, m - 1)$.
 - i. $m - 1 \geq kp$ and $(k + t_1 - 2) + k(k + s_1 - 2) \leq p$.
 - ii. $kp > m$ and $(k + t_1 - 2) + k(k + s_1 - 2) \leq p - b_j(n - k - s_1)$.

Definition 1.1 Let T be the set of triplets $(k, m, n) \in \mathbb{N}^3$ satisfying $k \geq 2, n > m \geq 2$ and $k + m^2 + n^2 - 2 > kmn$. For $(k, m, n) \in T$ there exists a $j \in \mathbb{N}$ such that $(\mathrm{GL}_1^k \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1 \otimes \omega_1)^{\oplus k})$ is transformed to a trivial prehomogeneous module by j castling transformations. This number j is uniquely determined if we use only castling transformations decreasing the module's dimension. This unique j will be denoted by $\nu(k, m, n)$. Thus we obtain a map $\nu : T \rightarrow \mathbb{N}$. For example, $\nu(k, m, n) = 0$ if and only if $mk \leq n$. We define (a_i) to be the sequence

$$a_{-1} = -1, \quad a_0 = 0, \quad a_i = ka_{i-1} - a_{i-2} \text{ for } i > 0.$$

There are some cases of the form KII IV belonging neither to KII-16 nor KII-17, but to KII I instead. These are the cases (4.1-i), (4.1-ii), (4.7) and (4.8) from Section 4.2 in Kimura et al. [5]. We will list them here for the sake of completeness.

1. $(\mathrm{GL}_1^{1+s+t} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1 \otimes 1)^{(*)\oplus s} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes (\omega_1^* \oplus \omega_1^{\oplus t-1})))$
with $k = 1, t \geq 1, n \geq m + t - 1$ and $s \leq m$. This is the case KII I-3.
2. $(\mathrm{GL}_1^{k+s+t} \times \mathrm{SL}_m \times \mathrm{SL}_n, ((\omega_1^{\oplus s-1} \oplus \omega_1^{(*)}) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1^* \oplus \omega_1^{\oplus t-1})))$
with $k \geq 2, t \geq 1, n \geq km + t - 1$ and $s + k \leq m + 1$. This is the case KII I-3.
3. $(\mathrm{GL}_1^{k+s+t} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1^{(*)} \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$
with $n \geq km + t$, and $(\mathrm{GL}_1^s \times \mathrm{SL}_m, \omega_1^{(*)\oplus s})$ is a simple prehomogeneous module. This is the case KII I-1.
4. $(\mathrm{GL}_1^{k+s+t} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1^{(*)} \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$
with $t \geq 3, n = km + t - 1$, and $(\mathrm{GL}_1^{k+s} \times \mathrm{SL}_m, \omega_1^{*\oplus k} \oplus \omega_1^{(*)\oplus s})$ is a simple prehomogeneous module. This is the case KII I-2.

2 Tables of étale modules

In this chapter, we present the étale modules from part IV of Globke [1]. Note that this is not claimed to be a complete classification.

2.1 Étale modules with center GL_1

2.1.1 One-simple étale modules with center GL_1

- SK I-4: $(\mathrm{GL}_2, 3\omega_1, \mathrm{Sym}^3 \mathbb{C}^2)$.
- Ks I-2: $(\mathrm{GL}_1 \times \mathrm{SL}_n, \mu \otimes \omega_1^{\oplus n}, (\mathbb{C}^n)^{\oplus n})$.

2.1.2 Two-simple étale modules with center GL_1

- SK I-8: $(\mathrm{SL}_3 \times \mathrm{GL}_2, 2\omega_1 \otimes \omega_1, \mathrm{Sym}^2 \mathbb{C}^3 \otimes \mathbb{C}^2)$.
- SK I-11: $(\mathrm{SL}_5 \times \mathrm{GL}_4, \omega_2 \otimes \omega_1, \bigwedge^2 \mathbb{C}^5 \otimes \mathbb{C}^4)$.
- KII I-2: $(\mathrm{GL}_1 \times G \times \mathrm{SL}_n, (\varrho_1 \otimes \omega_1) \oplus \dots \oplus (\varrho_k \otimes \omega_1) \oplus (\varrho_{k+1}^* \otimes 1) \oplus \dots \oplus (\varrho_l^* \otimes 1))$, with $n = -1 + \sum_{i=1}^k \dim(\varrho_i)$ and $(\mathrm{GL}_1 \times G, \varrho_1 \oplus \dots \oplus \varrho_l)$ an étale module for a simple group G .
- KII IV-16-iv (a): $(\mathrm{GL}_m \times \mathrm{SL}_{m+1}, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*)^{\oplus m+1}), m \geq 2$.

2.2 Étale modules for groups with factor Sp_m

- KI I-16: $(\mathrm{GL}_1^2 \times \mathrm{Sp}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1^*), (\mathbb{C}^4 \otimes \mathbb{C}^3) \oplus V^5 \oplus \mathbb{C}^3)$.
- KI I-18: $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1), (V^5 \otimes \mathbb{C}^2) \oplus \mathbb{C}^4 \oplus \mathbb{C}^2)$.
- KI I-19: $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_4, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*), (V^5 \otimes \mathbb{C}^4) \oplus \mathbb{C}^4 \oplus \mathbb{C}^4)$.

2.3 One-simple étale modules

- SK I-4: $(\mathrm{GL}_2, 3\omega_1, \mathrm{Sym}^3 \mathbb{C}^2)$.
- Ks I-2: $(\mathrm{GL}_1 \times \mathrm{SL}_n, \mu \otimes \omega_1^{\oplus n}, (\mathbb{C}^n)^{\oplus n})$.
- Ks I-3: $(\mathrm{GL}_1^{n+1} \times \mathrm{SL}_n, \omega_1^{\oplus n+1}, (\mathbb{C}^n)^{\oplus n+1})$.
- Ks I-4: $(\mathrm{GL}_1^{n+1} \times \mathrm{SL}_n, \omega_1^{\oplus n} \oplus \omega_1^*, (\mathbb{C}^n)^{\oplus n} \oplus \mathbb{C}^{n*})$.
- Ks I-11 for $n = 2$: $(\mathrm{GL}_1^2 \times \mathrm{SL}_2, 2\omega_1 \oplus \omega_1, \mathrm{Sym}^2 \mathbb{C}^2 \otimes \mathbb{C}^2)$.

2.4 Two-simple étale modules

- SK I-8: $(\mathrm{SL}_3 \times \mathrm{GL}_2, 2\omega_1 \otimes \omega_1, \mathrm{Sym}^2 \mathbb{C}^3 \otimes \mathbb{C}^2)$.
- SK I-11: $(\mathrm{SL}_5 \times \mathrm{GL}_4, \omega_2 \otimes \omega_1, \bigwedge^2 \mathbb{C}^5 \otimes \mathbb{C}^4)$.
- KI I-1: $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes \omega_1), (\bigwedge^2 \mathbb{C}^4 \otimes \mathbb{C}^2) \oplus (\mathbb{C}^4 \otimes \mathbb{C}^2))$.
- KI I-2: $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (\omega_1 \otimes 1), (\bigwedge^2 \mathbb{C}^4 \otimes \mathbb{C}^2) \oplus \mathbb{C}^4 \oplus \mathbb{C}^4)$.
- KI I-6: $(\mathrm{GL}_1^3 \times \mathrm{SL}_5 \times \mathrm{SL}_2, (\omega_2 \otimes \omega_1) \oplus (\omega_1^* \otimes 1) \oplus (\omega_1^{(*)} \otimes 1), (\bigwedge^2 \mathbb{C}^5 \otimes \mathbb{C}^2) \oplus \mathbb{C}^{5*} \oplus \mathbb{C}^{5(*)})$.

- KI I-16: $(\mathrm{GL}_1^2 \times \mathrm{Sp}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1^*), (\mathbb{C}^4 \otimes \mathbb{C}^3) \oplus V^5 \oplus \mathbb{C}^3)$.
- KI I-18: $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_2, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1), (\mathbb{C}^4 \otimes \mathbb{C}^2) \oplus \mathbb{C}^4 \oplus \mathbb{C}^2)$.
- KI I-19: $(\mathrm{GL}_1^3 \times \mathrm{Sp}_2 \times \mathrm{SL}_4, (\omega_1 \otimes \omega_1) \oplus (\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^*), (\mathbb{C}^4 \otimes \mathbb{C}^4) \oplus \mathbb{C}^4 \oplus \mathbb{C}^4)$.
- KII I-1: $(\mathrm{GL}_1^j \times G \times \mathrm{GL}_n, ((\sigma_1 \oplus \dots \oplus \sigma_s) \otimes \omega_1) \oplus ((\varrho_1 \oplus \dots \oplus \varrho_l) \otimes 1))$,
with $n = \sum_{i=1}^s \dim(\varrho_i)$ and $(\mathrm{GL}_1^j \times G, \varrho_1 \oplus \dots \oplus \varrho_l)$ an étale module for a simple group G , $1 \leq j \leq l$.
- KII I-2: $(\mathrm{GL}_1^{j+t} \times G \times \mathrm{SL}_n, ((\varrho_1 \oplus \dots \oplus \varrho_k) \otimes \omega_1) \oplus ((\varrho_{k+1}^* \oplus \dots \oplus \varrho_l^*) \otimes 1) \oplus (1 \otimes \omega_1^{\oplus t}))$,
with $n = t - 1 + \sum_{i=1}^k \dim(\varrho_i)$, $1 \leq j \leq l$, and $(\mathrm{GL}_1^j \times G, \varrho_1 \oplus \dots \oplus \varrho_l)$ an étale module for a simple group G .
- KII I-3: $(\mathrm{GL}_1^{j+t} \times G \times \mathrm{SL}_n, ((\varrho_1 \oplus \dots \oplus \varrho_k) \otimes \omega_1) \oplus ((\varrho_{k+1} \oplus \dots \oplus \varrho_l) \otimes 1) \oplus (1 \otimes (\omega_1^{\oplus t-1} \oplus \omega_1^*)))$,
with $n = t - 1 + \sum_{i=1}^k \dim(\varrho_i)$, $1 \leq j \leq l$, and $(\mathrm{GL}_1^j \times G, \varrho_1 \oplus \dots \oplus \varrho_l)$ an étale module for a simple group G .
- KII II-4-i (b): $(\mathrm{GL}_1^3 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes 2\omega_1^{(*)}) \oplus (\omega_1^{(*)} \otimes 1))$.
- KII II-4-ii (a): $(\mathrm{GL}_1^5 \times \mathrm{SL}_3 \times \mathrm{SL}_4, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus ((\omega_1 \oplus \omega_1)^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$.
- KII II-4-iii (d): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus ((\omega_1 \oplus \omega_1)^{(*)} \otimes 1) \oplus (1 \otimes \omega_1))$.
- KII II-4-iii (f): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1))$.
- KII II-4-iii (g): $(\mathrm{GL}_1^4 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*))$.
- KII II-4-iii (h): $(\mathrm{GL}_1^4 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*))$.
- KII II-4-iii (n): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus ((\omega_1 \oplus \omega_1)^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$.
- KII II-4-iii (p): $(\mathrm{GL}_1^4 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*))$.
- KII II-4-iii (q): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (\omega_1 \otimes 1) \oplus (\omega_1^* \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$.
- KII II-5-i (b): $(\mathrm{GL}_1^3 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (2\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^{(*)}) \oplus (\omega_1 \otimes 1))$.
- KII II-5-ii (b): $(\mathrm{GL}_1^4 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (2\omega_1 \otimes 1) \oplus (1 \otimes \omega_1))$.
- KII II-5-iii (e): $(\mathrm{GL}_1^4 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (2\omega_1 \otimes 1) \oplus (1 \otimes \omega_1^{(*)}))$.
- KII II-5-iv (a): $(\mathrm{GL}_1^3 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_2^*))$.
- KII II-6 (b): $(\mathrm{GL}_1^3 \times \mathrm{SL}_3 \times \mathrm{SL}_5, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2) \oplus (1 \otimes \omega_2))$.
- KII II-9 (a): $(\mathrm{GL}_1^4 \times \mathrm{SL}_6 \times \mathrm{SL}_7, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_2^*) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1))$.
- KII II-10: $(\mathrm{GL}_1^{r+1} \times \mathrm{SL}_n \times \mathrm{SL}_n, (\omega_1 \otimes \omega_1) \oplus (\varrho_1 \otimes 1) \oplus \dots \oplus (\varrho_k \otimes 1) \oplus (1 \otimes \varrho_{k+1}^*) \oplus \dots \oplus (1 \otimes \varrho_r^*))$, where $(\mathrm{GL}_1^r \times \mathrm{SL}_n, \varrho_1 \oplus \dots \oplus \varrho_r)$ is an étale module.
- KII III-12: $(\mathrm{GL}_1^2 \times \mathrm{SL}_4 \times \mathrm{SL}_8, (\omega_2 \otimes \omega_1) \oplus (\omega_1 \otimes \omega_1))$.
- KII III-13-i (d): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^{(*)} \otimes 1) \oplus (1 \otimes \omega_1^*))$.

- KII III-13-i (e): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2^* \otimes 1) \oplus (1 \otimes \omega_1))$.
- KII III-13-i (g): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*) \oplus (1 \otimes \omega_1^*) \oplus (\omega_2 \otimes 1) \oplus (1 \otimes \omega_1))$.
- KII III-13-ii (a): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\omega_2 \otimes 1))$.
- KII III-13-ii (b): $(\mathrm{GL}_1^5 \times \mathrm{SL}_2 \times \mathrm{SL}_3, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\omega_2^* \otimes 1))$.
- KII IV-16-i: $(\mathrm{GL}_1^{n+2} \times \mathrm{SL}_n \times \mathrm{SL}_n, (\omega_1 \otimes 1)^{\oplus s_1} \oplus (\omega_1^* \otimes 1)^{\oplus s_2} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1)^{\oplus t_1} \oplus (1 \otimes \omega_1^*)^{\oplus t_2})$,
with $s_1 + t_2 = n$ and $s_2 + t_1 = 1$.
- KII IV-16-i: $(\mathrm{GL}_1^{n+2} \times \mathrm{SL}_n \times \mathrm{SL}_n, (\omega_1 \otimes 1)^{\oplus s_1} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*)^{\oplus t_2})$,
with $s_1 + t_2 = n + 1$.
- KII IV-16-i: $(\mathrm{GL}_n \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s_1} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*)^{\oplus t_2})$,
with $s_1 + t_2 = n$.
- KII IV-16-ii (a): $(\mathrm{GL}_n \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s_1} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1^*)^{\oplus t_2})$,
with $s_1 + kt_2 = m$.
- KII IV-16-ii (b): $(\mathrm{GL}_n \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s_2} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1)^{\oplus t_1})$,
with $s_2 + kt_1 = m$.
- KII IV-16-iii: $(\mathrm{GL}_1^{t_1-1} \times \mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s_2} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\mu \otimes 1 \otimes \omega_1)^{\oplus t_1-1})$,
with $s_2 + t_1 = m + 1$.
- KII IV-16-iv (a): $(\mathrm{GL}_1^{m+1} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\mu \otimes \omega_1^* \otimes 1) \oplus (\omega_1 \otimes \omega_1) \oplus (\mu \otimes 1 \otimes \omega_1^*)^{\oplus m})$,
with $n = m + 1$.
- KII IV-16-iv (a): $(\mathrm{GL}_m \times \mathrm{SL}_n, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*)^{\oplus m})$,
with $n = m + 1$.
- KII IV-16-iv (a): $(\mathrm{GL}_1^{m+2} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1^*)^{\oplus m+1})$,
with $n = m + 1$.
- KII IV-16-iv (a): $(\mathrm{GL}_1^{m+1} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\mu \otimes \omega_1 \otimes 1) \oplus (\omega_1 \otimes \omega_1) \oplus (\mu \otimes 1 \otimes \omega_1^*)^{\oplus m})$,
with $n = m + 1$.
- KII IV-16-iv (c): $(\mathrm{GL}_1^{t_2} \times \mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s_1} \oplus (\omega_1 \otimes \omega_1) \oplus (1 \otimes \omega_1) \oplus (\mu \otimes 1 \otimes \omega_1^*)^{\oplus t_2})$,
with $s_1 + t_2 = m$ and $n = m + 1$.
- KII IV-17 (1a): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1^*)^{\oplus t})$,
with $sa_{j+1} + ta_{j+2} = m_0$ and $n_0 = km_0$.

Let $p = km + t - n$ and $q = kp - m = k^2m + kt - kn - m$.

- KII IV-17 (1b): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp > m$, $s > 0$, $ta_{j+1} + sa_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$ for $\tilde{m}_0 = a_{j+1}p - a_jm$,
 $\tilde{n}_0 = a_jp - a_{j-1}m$ and $j = \nu(k, p, m)$.
- KII IV-17 (1b): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp > m$, $kq = p$, $2 \leq t$ and $kt = q$.
- KII IV-17 (1b): $(\mathrm{GL}_1^{k+1} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1))$,
with $kp > m$, $k = q + 1$ and $p = q^2 + q$.

- KII IV-17 (1b): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1))$,
with $kp > m$, $k = q$ and $p = q^2$.
- KII IV-17 (1b): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp > m$, $kq > p$, $ta_{j+2} = \tilde{m}_0$ and $k\tilde{m}_0 = \tilde{n}_0$ for $\tilde{m}_0 = a_{j+1}q - a_jp$, $\tilde{n}_0 = a_jq - a_{j-1}p$ and $j = \nu(k, q, p)$.
- KII IV-17 (1b): $(\mathrm{GL}_1^{1+k+t} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1 \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp = m$ and $t + k = p + 1$.
- KII IV-17 (1b): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp = m$ and $t + k = p$.
- KII IV-17 (1b): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp = m$, $2 \leq s$ and $t + ks = p$.

Let $p = km + t - n$, $q = kp + s - m$ and $r = kq - p$.

- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp > m$, $kq > p$, $t > 0$, $sa_{j+1} + ta_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$ for $\tilde{m}_0 = a_{j+1}q - a_jp$,
 $\tilde{n}_0 = a_jq - a_{j-1}p$ and $j = \nu(k, q, p)$.
- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $kp > m$, $kq > p$, $kr = q$, $2 \leq s$ and $ks = r$.
- KII IV-17 (1c): $(\mathrm{GL}_1^{k+1} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1^* \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $kp > m$, $kq > p$, $k = r + 1$ and $q = r^2 + r$.
- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $kp > m$, $kq > p$, $k = r$ and $q = r^2$.
- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $kp > m$, $kq > p$, $kr > q$, $sa_{j+2} = \tilde{m}_0$ and $k\tilde{m}_0 = \tilde{n}_0$ for $\tilde{m}_0 = a_{j+1}r - a_jq$,
 $\tilde{n}_0 = a_jr - a_{j-1}q$ and $j = \nu(k, r, q)$.
- KII IV-17 (1c): $(\mathrm{GL}_1^{s+k+1} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1))$,
with $kp > m$, $kq = p$ and $s + k = q + 1$.
- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1))$,
with $kp > m$, $kq = p$ and $s + k = q$.
- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1)^{\oplus t})$,
with $kp > m$, $kq = p$, $2 \leq t$ and $s + kt = q$.
- KII IV-17 (1c): $(\mathrm{GL}_1^s \times \mathrm{SL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $k = p$ and $kp + s - 1 = m$.
- KII IV-17 (1c): $(\mathrm{GL}_1^{s+k} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $k = p + 1$ and $kp + s - 1 = m$.
- KII IV-17 (1c): $(\mathrm{GL}_1^{s+k+1} \times \mathrm{SL}_m \times \mathrm{SL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1))$,
with $k = p$ and $kp + s - 1 = m$.
- KII IV-17 (1c): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1^* \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k})$,
with $ks = p$ and $kp = m$.

Let $p = km + t - n - 1$.

- KII IV-17 (2a): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1^* \oplus \omega_1^{\oplus t-1})))$,
with $kp > m$, $(t-2)a_{j+1} + (k+s)a_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$ for $\tilde{m}_0 = a_{j+1}p - a_jm$,
 $\tilde{n}_0 = a_jp - a_{j-1}m$ and $j = \nu(k, p, m)$.
- KII IV-17 (2a): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1^* \oplus \omega_1^{\oplus t-1})))$,
with $kp = m$ and $t-2+k(k+s) = p$.

Let $p = km + t - n$ and $q = kp + t - m - 1$.

- KII IV-17 (2b): $(\mathrm{GL}_m \times \mathrm{GL}_n, ((\omega_1 \oplus \omega_1^{*\oplus s-1}) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1^{\oplus t}))$,
with $kp > m$, $kq > p$, $(s-2)a_{j+1} + (k+t)a_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$ for $\tilde{m}_0 = a_{j+1}q - a_jp$, $\tilde{n}_0 = a_jq - a_{j-1}p$ and $j = \nu(k, q, p)$.
- KII IV-17 (2b): $(\mathrm{GL}_m \times \mathrm{GL}_n, ((\omega_1 \oplus \omega_1^{*\oplus s-1}) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1^{\oplus t}))$,
with $kp > m$, $kq = p$ and $s-2+k(k+t) = q$.
- KII IV-17 (2b): $(\mathrm{GL}_1^{k+s+t} \times \mathrm{SL}_m \times \mathrm{SL}_n, ((\omega_1 \oplus \omega_1^{*\oplus s-1}) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1^{\oplus t}))$,
with $m = s-1+kp$ and $k+t = p+1$.
- KII IV-17 (2b): $(\mathrm{GL}_1^s \times \mathrm{SL}_m \times \mathrm{GL}_n, ((\omega_1 \oplus \omega_1^{*\oplus s-1}) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes \omega_1^{\oplus t}))$,
with $m = s-1+kp$ and $k+t = p$.
- KII IV-17 (3a): $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1 \oplus \omega_1^{*\oplus t-1})))$,
with $(k+s)a_{j+1} + (t-2)a_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$, for $\tilde{m}_0 = a_{j+1}m - a_j(n-1)$,
 $\tilde{n}_0 = a_jm - a_{j-1}(n-1)$ and $j = \nu(k, m, n-1)$.

Let $p = km + t - n$.

- KII IV-17 (3b) $(\mathrm{GL}_m \times \mathrm{GL}_n, (\omega_1 \otimes 1)^{\oplus s} \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1 \oplus \omega_1^{*\oplus t-1})))$,
with $kp > m$, $(k+t)a_{j+1} + (s-2)a_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$, for $\tilde{m}_0 = a_{j+1}p - a_j(m-1)$,
 $\tilde{n}_0 = a_jp - a_{j-1}(m-1)$ and $j = \nu(k, p, m-1)$.

Let $p = km + t - n - 1$.

- KII IV-17 (4): $(\mathrm{GL}_m \times \mathrm{GL}_n, ((\omega_1^{\oplus s-1} \oplus \omega_1^*) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1^{\oplus t-1} \oplus \omega_1^*)))$,
with $kp > m-1$, $(k+t-2)a_{j+1} + (k+s-2)a_{j+2} = \tilde{m}_0$ and $\tilde{n}_0 = k\tilde{m}_0$ for
 $\tilde{m}_0 = a_{j+1}p - a_j(m-1)$, $\tilde{n}_0 = a_jp - a_{j-1}(m-1)$ and $j = \nu(k, p, m-1)$.
- KII IV-17 (4): $(\mathrm{GL}_m \times \mathrm{GL}_n, ((\omega_1^{\oplus s-1} \oplus \omega_1^*) \otimes 1) \oplus (\omega_1 \otimes \omega_1)^{\oplus k} \oplus (1 \otimes (\omega_1^{\oplus t-1} \oplus \omega_1^*)))$,
with $m-1 = kp$ and $(k+t-2) + k(k+s-2) = p$.

References

- [1] W. GLOBKE
Étale Representations of Reductive Algebraic Groups and Left-Symmetric Algebras
Diploma thesis, University of Karlsruhe, 2007
- [2] T. KIMURA
Introduction to Prehomogeneous Vector Spaces
American Mathematical Society, 2003
- [3] T. KIMURA
A Classification of Prehomogeneous Vector Spaces of Simple Algebraic Groups with Scalar Multiplications
J. Algebra 83, 1983, 72-100
- [4] T. KIMURA, S. KASAI, M. INUZUKA, O. YASUKURA
A Classification of 2-Simple Prehomogeneous Vector Spaces of Type I
J. Algebra 114, 1988, 369-400
- [5] T. KIMURA, S. KASAI, M. TAGUCHI, M. INUZUKA
Some P.V.-Equivalences and a Classification of 2-Simple Prehomogeneous Vector Spaces of Type II
Trans. AMS 308, 2, 1988, 433-494
- [6] T. KIMURA, T. KOGISO, K. SUGIYAMA
Relative Invariants of 2-Simple Prehomogeneous Vector Spaces of Type I
J. Algebra 308, 2007, 445-483
- [7] T. KOGISO, G. MIYABE, M. KOBAYASHI, T. KIMURA
Nonregular 2-Simple Prehomogeneous Vector Spaces of Type I and Their Relative Invariants
J. Algebra 251, 2002, 27-69
- [8] M. SATO, T. KIMURA
A Classification of Irreducible Prehomogeneous Vector Spaces and their Relative Invariants
Nagoya Math. J. 65, 1977, 1-155